

Patient information

Call or see your oncologist or get emergency help RIGHT AWAY if you have any of these symptoms:

Fever (100°F/38°C or higher)	Severe fatigue or weakness
Fast or irregular heartbeat	Dizziness or unsteadiness
Difficulty finding words	Difficulty breathing
Chills or shaking chills	Seizures
Confusion	Tremors

Patient card

Cytokine Release Syndrome is a group of symptoms caused by small proteins called cytokines that are released in your body during inflammation.

SHOW THIS CARD if you go to the emergency room or see any physician.

This patient has received a T cell immunotherapy for multiple myeloma.

Caregiver information

Call or see your oncologist or get emergency help RIGHT AWAY if you see any of these symptoms:

Fever (100°F/38°C or higher)	Severe fatigue or weakness
Fast or irregular heartbeat	Dizziness or unsteadiness
Difficulty finding words	Difficulty breathing
Chills or shaking chills	Seizures
Confusion	Tremors

Caregiver card

Cytokine Release Syndrome is a group of symptoms caused by small proteins called cytokines that are released in your body during inflammation.

SHOW THIS CARD if the patient goes to the emergency room or sees any physician.

This patient has received a T cell immunotherapy for multiple myeloma.

Important Information for Physicians

GO39775

Name of treating oncologist:

This patient has received a T cell immunotherapy.

Office phone:

Drug can cause Cytokine Release Syndrome (CRS) and neurologic toxicities, which may be fatal or life threatening. CRS may involve any organ system.

After-hours phone:

It can occur in the 24-48 hours following infusion.

Date of infusion:

For physicians: References for CRS management can be found here:

www.edmesafe.com/bfcr4350a

Contact their oncologist immediately for further info

Important Information for Physicians

GO39775

Name of treating oncologist:

This patient has received a T cell immunotherapy.

Office phone:

Drug can cause Cytokine Release Syndrome (CRS) and neurologic toxicities, which may be fatal or life threatening. CRS may involve any organ system.

After-hours phone:

It can occur in the 24-48 hours following infusion.

Date of infusion:

For physicians: References for CRS management can be found here:

www.edmesafe.com/bfcr4350a

Contact their oncologist immediately for further info